

Asymptotic equivalence for nonparametric regression with multivariate and random design

Markus Reiß

Institute of Applied Mathematics

University of Heidelberg

reiss@statlab.uni-heidelberg.de

February 2, 2008

Abstract

We show that nonparametric regression is asymptotically equivalent in Le Cam's sense with a sequence of Gaussian white noise experiments as the number of observations tends to infinity. We propose a general constructive framework based on approximation spaces, which permits to achieve asymptotic equivalence even in the cases of multivariate and random design.

Key words and Phrases: Le Cam deficiency, equivalence of experiments, approximation space, interpolation, Gaussian white noise.

AMS subject classification: 62G08, 62G20, 62B15

1 Introduction

Nonparametric regression is the model most often encountered in nonparametric statistics because of its widespread applications. On the other hand, for theoretical investigations the Gaussian white noise or sequence space model is often preferred since it exhibits nice mathematical properties. The common wisdom that statistical decisions in the two models show the same asymptotic behaviour has been formalized and proved for the first time by Brown and Low (1996) in the one-dimensional case, using Le Cam's concept of equivalence of statistical experiments.

In this paper we propose a unifying framework for establishing global asymptotic equivalence between Gaussian nonparametric regression and white noise experiments based on constructive transitions with only minimal randomisations. This framework not only allows to give concise proofs of known results, but extends the asymptotic equivalence to the multivariate and random design situation. The multivariate result has often been alluded to, though it has never been proved, see e.g. Hoffmann and Lepski (2002). While Brown and Zhang (1998) remark that the regression and white noise experiments are not asymptotically equivalent for equidistant design on $[0, 1]^d$ and Sobolev classes of regularity $s \leq d/2$, the so far only positive result by Carter (2006) states asymptotic equivalence for equidistant design in dimensions $d = 2$ and $d = 3$ when $s > d/2$. The difficulty in extending results to higher dimensions is that we have to go beyond piecewise constant or linear approximations. For the dynamic model of ergodic diffusions Dalalyan and Reiß (2006) have established multidimensional asymptotic equivalence with a white noise model. For the case of univariate nonparametric regression, but with non-Gaussian errors we refer to Grama and Nussbaum (1998).

In Section 2 the concept of isometric approximation spaces is introduced and applied to local constant and Fourier approximations. The latter yields an easy proof for asymptotic equivalence in any dimension d for periodic Sobolev classes of regularity $s > d/2$ and extends scalar results by Rohde (2004). A more flexible

framework is obtained using isomorphic approximations spaces in Section 3. As a main application, a constructive asymptotic equivalence result is established on the basis of wavelet multiresolution analyses, which provides equivalence results also for non-periodic function classes. Connections to asymptotic studies by Donoho and Johnstone (1999) and Johnstone and Silverman (2004) for wavelet estimators are discussed. The case of a random design, uniform on a d -dimensional cube, is treated in Section 4. This setting is much more involved, but can also be cast in the isomorphic framework. The construction is based on a two-level procedure, generalizing an idea by Brown, Cai, Low, and Zhang (2002). Fine approximation and symmetry properties of the Fourier basis yield the main result that also in the case of random design asymptotic equivalence holds for Sobolev regularities $s > d/2$ and any dimension $d \geq 1$.

2 Isometric approximation

2.1 General theory

We write $\mathcal{L}^2(\mathcal{D}) := \{f : \mathcal{D} \rightarrow \mathbb{K} \mid \|f\|_{L^2}^2 := \int |f|^2 < \infty\}$ with $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$ and $L^2(\mathcal{D})$ for the Hilbert space of equivalence classes with respect to $\|\cdot\|_{L^2}$. Although the observations are real-valued, we shall use complex-valued functions for simplicity when treating Fourier approximations.

2.1 Definition. Let $\mathbb{E}_n^d$ be the regression experiment obtained from observing

$$Y_i = f(x_i) + \sigma \varepsilon_i, \quad i = 1, \dots, n,$$

for $n \in \mathbb{N}$, $f : \mathcal{D} \subseteq \mathbb{R}^d \rightarrow \mathbb{R}$ in some class $\mathcal{F}^d \subseteq \mathcal{L}^2(\mathcal{D})$, fixed design points $x_i \in \mathcal{D}$ and for independent random variables $\varepsilon_i \sim \mathcal{N}(0, 1)$.

Suppose we are given an n -dimensional space $S_n \subseteq \mathcal{L}^2(\mathcal{D})$ and a linear mapping $D_n : \mathcal{L}^2(\mathcal{D}) \rightarrow \mathbb{K}^n$ with the following isometric property on S_n

$$\forall g_n \in S_n : \|g_n\|_{L^2} = \|g_n\|_n := n^{-1/2} \|D_n g_n\|_{\mathbb{K}^n}. \quad (2.1)$$

By $\langle \bullet, \bullet \rangle_n$ we denote the scalar product associated with $\|\bullet\|_n$. Usually, $D_n g = (g(x_i))_{1 \leq i \leq n}$ will be the point evaluation at the n design points in which case $\|g\|_n^2 = \frac{1}{n} \sum_{i=1}^n |g(x_i)|^2$ is just the empirical norm. Let us further introduce the linear operator

$$\mathcal{J}_n : \mathcal{L}^2(\mathcal{D}) \rightarrow S_n, \quad \mathcal{J}_n g := (D_n|_{S_n})^{-1} (g(x_1), \dots, g(x_n))^\top.$$

For $D_n g = (g(x_i))_{1 \leq i \leq n}$ we have $\mathcal{J}_n = (D_n|_{S_n})^{-1} D_n$ and $\mathcal{J}_n$ is the $\|\bullet\|_n$ -orthogonal projection onto S_n such that $\mathcal{J}_n g$ is the unique element of S_n interpolating g at the design points (x_i) .

To state the first results, we refer to Le Cam and Lo Yang (2000) for the notion of equivalence between experiments and of the Le Cam distance between two experiments $\mathbb{E}$ and $\mathbb{G}$, which for the parameter class $\mathcal{F}$ will be denoted by $\Delta_{\mathcal{F}}(\mathbb{E}, \mathbb{G})$. The Gaussian law on a Hilbert space H with mean vector $\mu \in H$ and covariance operator $\Sigma : H \rightarrow H$ will be denoted by $\mathcal{N}(\mu, \Sigma)$.

The regression experiment $\mathbb{E}_n^d$ can be transformed to a functional Gaussian shift experiment by applying the isometry $(D_n|_{S_n})^{-1}$ to $Y = (Y_i) \in \mathbb{R}^n$:

$$Z := (D_n|_{S_n})^{-1} Y = \mathcal{J}_n f + \frac{\sigma}{\sqrt{n}} \zeta \in S_n, \quad (2.2)$$

where $\zeta := \sqrt{n}(D_n|_{S_n})^{-1} \varepsilon \sim \mathcal{N}(0, \text{Id}_{S_n})$ is a Gaussian white noise in S_n because for $g_n, h_n \in S_n$

$$E[\langle \zeta, g_n \rangle_{L^2} \overline{\langle \zeta, h_n \rangle_{L^2}}] = n^{-1} E[\langle \varepsilon, D_n g_n \rangle_{\mathbb{K}^n} \overline{\langle \varepsilon, D_n h_n \rangle_{\mathbb{K}^n}}] = \langle g_n, h_n \rangle_n = \langle g_n, h_n \rangle_{L^2}.$$

By adding completely uninformative observations on the orthogonal complement of S_n in $L^2(\mathcal{D})$, the observation of Z in (2.2) is equivalent to observing

$$\langle \varphi, Z \rangle_{L^2} = \langle \varphi, \mathcal{J}_n f \rangle_{L^2} + \frac{\sigma}{\sqrt{n}} \langle \varphi, \bar{\zeta} \rangle_{L^2} \quad \forall \varphi \in L^2(\mathcal{D}),$$

with $\langle \varphi, \bar{\zeta} \rangle_{L^2} \sim \mathcal{N}(0, \|\varphi\|_{L^2}^2)$. In differential notation we have thus established the following equivalence.

2.2 Proposition. *Let $\mathbb{F}_n^d$ be the Gaussian white noise experiment in $L^2(\mathcal{D})$ given by observing*

$$dY(x) = \mathcal{I}_n f(x) dx + \frac{\sigma}{\sqrt{n}} dB(x), \quad x \in \mathcal{D},$$

where $f \in \mathcal{F}^d$ and dB is a Gaussian white noise in $L^2(\mathcal{D})$. Then the regression experiment $\mathbb{E}_n^d$ is statistically equivalent to $\mathbb{F}_n^d$ for any functional class $\mathcal{F}^d$.

We are coming to the first main result.

2.3 Definition. *Let $\mathbb{G}_n^d$ be the Gaussian white noise experiment given by observing*

$$dY(x) = f(x) dx + \frac{\sigma}{\sqrt{n}} dB(x), \quad x \in \mathcal{D},$$

where $f \in \mathcal{F}^d$ and dB is a Gaussian white noise in $L^2(\mathcal{D})$.

2.4 Theorem. *The Le Cam distance between $\mathbb{E}_n^d$ and $\mathbb{G}_n^d$ for the class $\mathcal{F}^d$ is bounded by*

$$\Delta_{\mathcal{F}^d}(\mathbb{E}_n^d, \mathbb{G}_n^d) \leq 1 - 2\Phi\left(-\frac{\sqrt{n}}{2\sigma} \sup_{f \in \mathcal{F}^d} \|f - \mathcal{I}_n f\|_{L^2}\right),$$

where Φ denotes the standard Gaussian cumulative distribution function.

2.5 Remark. *Note that $\|f - \mathcal{I}_n f\|_{L^2}^2 = \|f - P_n f\|_{L^2}^2 + \|P_n f - \mathcal{I}_n f\|_{L^2}^2$ holds where P_n is the L^2 -orthogonal projection onto S_n . This means that the bound on the Le Cam distance is always larger than the same expression involving the classical bias estimate $\sup_{f \in \mathcal{F}^d} \|f - P_n f\|_{L^2}$. Because of $\Phi(0) = 1/2$ Proposition 2.4 yields the rate estimate*

$$\Delta_{\mathcal{F}^d}(\mathbb{E}_n^d, \mathbb{G}_n^d) \lesssim \sigma^{-1} n^{1/2} \sup_{f \in \mathcal{F}^d} \|f - \mathcal{I}_n f\|_{L^2}.$$

Here and in the sequel $A \lesssim B$ means $A \leq cB$ with a constant $c > 0$, independent of the other parameters involved, and $A \sim B$ is short for $A \lesssim B$ and $B \lesssim A$.

Proof. Since $\mathbb{E}_n^d$ and $\mathbb{F}_n^d$ are equivalent, it suffices to establish the bound for $\Delta_{\mathcal{F}^d}(\mathbb{F}_n^d, \mathbb{G}_n^d)$. The two latter experiments are realized on the same sample space. Therefore the Le Cam distance is bounded by the maximal total variation distance

over the class $\mathcal{F}^d$ (Nussbaum 1996, Prop. 2.2). For Gaussian white noise the total variation distance is given by $1 - 2\Phi(-\frac{\sqrt{n}}{2\sigma}\|f - \mathcal{J}_n f\|_{L^2})$ (Carter 2006, Section 3.2), and the result follows. $\square$

2.2 Piecewise constant approximation

The original results of Brown and Low (1996) for equidistant design on $\mathcal{D} = (0, 1]$ fit into the proposed isometric framework. For design points $x_i = i/n$, $i = 1, \dots, n$, we consider the n -dimensional space S_n of piecewise constant, left-continuous functions on $(0, 1]$ with possible jumps at i/n , $i = 1, \dots, n-1$. Using $D_n g = (g(i/n))_{1 \leq i \leq n}$ we obtain for $g_n \in S_n$

$$\|g_n\|_n^2 = \frac{1}{n} \sum_{i=1}^n |g_n(i/n)|^2 = \sum_{i=1}^n \int_{(i-1)/n}^n |g_n(u)|^2 du = \|g_n\|_{L^2}^2$$

such that D_n has the isometric property. To infer asymptotic equivalence by Proposition 2.4, we have to ensure that $\|f - \mathcal{J}_n f\|_{L^2} = o(n^{-1/2})$ uniformly over all f in some functional class $\mathcal{F}^d$. Considering the Hölder class of regularity $\alpha \in (0, 1]$

$$\mathcal{F}_H(\alpha, R) := \left\{ f \in C^\alpha([0, 1]) \mid \sup_{x \neq y} |f(x) - f(y)| / |x - y|^\alpha \leq R \right\},$$

we obtain for $f \in \mathcal{F}_H(\alpha, R)$

$$\begin{aligned} \|f - \mathcal{J}_n f\|_{L^2}^2 &= \sum_{i=1}^n \int_{(i-1)/n}^{i/n} |f(x) - f(i/n)|^2 dx \\ &\leq R^2 \sum_{i=1}^n \int_{(i-1)/n}^{i/n} |x - i/n|^{2\alpha} dx \\ &= R^2 (2\alpha + 1)^{-1} n^{-2\alpha}. \end{aligned}$$

Consequently, asymptotic equivalence between $\mathbb{E}_n^1$ and $\mathbb{G}_n^1$ holds for any Hölder class $\mathcal{F}_H(\alpha, R)$ with $\alpha > 1/2$ and $R > 0$ arbitrary. The approximation property of the Haar wavelet yields even asymptotic regularity for L^2 -Sobolev classes of regularity $\alpha > 1/2$.

For nonuniform design $0 \leq x_1 < \dots < x_n \leq 1$ consider the same setting as

before, in particular $D_n g = (g(i/n))_i \neq (g(x_i))_i$. We obtain for $f \in \mathcal{F}_H(\alpha, R)$:

$$\begin{aligned} \|f - \mathcal{J}_n f\|_{L^2}^2 &= \sum_{i=1}^n \int_{(i-1)/n}^{i/n} |f(x) - f(x_i)|^2 dx \leq R^2 \sum_{i=1}^n \int_{(i-1)/n}^{i/n} |x - x_i|^{2\alpha} dx \\ &\leq R^2 n^{-1} \sum_{i=1}^n \left(n^{-1} + |x_i - i/n| \right)^{2\alpha} \leq 2R^2 n^{-2\alpha} + 2R^2 n^{-1} \sum_{i=1}^n |x_i - i/n|^{2\alpha}. \end{aligned}$$

By Theorem 2.4 we have obtained the following result.

2.6 Theorem. *On the Hölder class $\mathcal{F}_H(\alpha, R)$ the Le Cam distance between nonparametric regression with design $0 < x_1^{(n)} < \dots < x_n^{(n)} \leq 1$ and the white noise experiment satisfies*

$$\Delta_{\mathcal{F}_H(\alpha, R)}(\mathbb{E}_n^1, \mathbb{G}_n^1) \lesssim \sigma^{-1} R \left(n^{1-2\alpha} + \sum_{i=1}^n |x_i^{(n)} - i/n|^{2\alpha} \right)^{1/2}.$$

Consequently, asymptotic equivalence holds whenever $\alpha \in (1/2, 1]$ and the design satisfies $\lim_{n \rightarrow \infty} \sum_{j=1}^n |x_j^{(n)} - i/n|^{2\alpha} = 0$, e.g. if $\max_i |x_i^{(n)} - i/n| = o(n^{-1/(2\alpha)})$.

2.7 Remark. *This approach does not permit to establish global equivalence for the random design case in Section 4 because the standard deviations of the order statistics $X_{(j)}$ decrease only with rate $n^{-1/2}$. Treating the random design like being equidistant yields nevertheless for estimation purposes nearly optimal asymptotic L^2 -risk when $\alpha > 1/2$ (Cai and Brown 1999).*

2.3 Fourier series approximation

In the case of $\mathcal{D} = [0, 1]^d$, $d \geq 1$, and of an equidistant design $(k/m)_{k \in \{1, \dots, m\}^d}$ with $m = n^{1/d} \in \mathbb{N}$ and odd, the Fourier system ($\iota := \sqrt{-1}$)

$$\varphi_\ell(x) := \exp(2\pi \iota \langle x, \ell \rangle), \quad \ell = (\ell_1, \dots, \ell_d), \quad |\ell|_\infty \leq \frac{m-1}{2},$$

is not only L^2 -orthonormal, but also with respect to $\langle \bullet, \bullet \rangle_n$ for $D_n g := (g(k/m))_k$:

$$\begin{aligned}
\langle \varphi_\ell, \varphi_{\ell'} \rangle_n &= \frac{1}{n} \sum_{k \in \{1, \dots, m\}^d} \varphi_\ell(k/m) \overline{\varphi_{\ell'}(k/m)} \\
&= m^{-d} \sum_{k_1, \dots, k_d=1}^m \prod_{i=1}^d \exp\left(2\pi i k_i (\ell_i - \ell'_i)/m\right) \\
&= \prod_{i=1}^d \left(\frac{1}{m} \sum_{\kappa=1}^m \exp\left(2\pi i \kappa (\ell_i - \ell'_i)/m\right) \right) \\
&= \begin{cases} 1, & \text{if } m | (\ell_i - \ell'_i) \text{ for all } i, \\ 0, & \text{otherwise.} \end{cases} \tag{2.3}
\end{aligned}$$

Consequently, the space of trigonometric polynomials $S_n := \text{span}(\varphi_\ell, |\ell|_\infty \leq \frac{m-1}{2})$ satisfies the isometric property (2.1).

The d -dimensional periodic Sobolev class of regularity s and radius R on $[0, 1]^d$ is given by

$$\mathcal{F}_{S, \text{per}}^d(s, R) := \left\{ f \in L^2([0, 1]^d) \mid \sum_{\ell \in \mathbb{Z}^d} |\ell|_\infty^{2s} |\langle f, \varphi_\ell \rangle|^2 \leq R^2 \right\}.$$

Due to the strong cancellation property (2.3) of the scalar product $\langle \bullet, \bullet \rangle_n$ we derive explicitly

$$(\mathcal{I}_n f)(x) = \sum_{|\ell|_\infty \leq (m-1)/2} \left(\sum_{k \in \mathbb{Z}^d} \langle f, \varphi_{\ell+km} \rangle \right) \varphi_\ell(x).$$

In view of Remark 2.5 we first bound the classical bias:

$$\sup_{f \in \mathcal{F}_{S, \text{per}}^d(s, R)} \|f - P_n f\|_{L^2}^2 = \sup_{f \in \mathcal{F}_{S, \text{per}}^d(s, R)} \sum_{|\ell|_\infty \geq (m+1)/2} |\langle f, \varphi_\ell \rangle|^2 = R^2 \left(\frac{m+1}{2} \right)^{-2s}.$$

For $s > d/2$ we obtain, using the Cauchy-Schwarz inequality,

$$\begin{aligned}
&\sup_{f \in \mathcal{F}_{S, \text{per}}^d(s, R)} \|P_n f - \mathcal{I}_n f\|_{L^2}^2 \\
&= \sup_{f \in \mathcal{F}_{S, \text{per}}^d(s, R)} \sum_{|\ell|_\infty \leq (m-1)/2} \left(\sum_{k \in \mathbb{Z}^d \setminus \{0\}} \langle f, \varphi_{\ell+km} \rangle \right)^2 \\
&\leq \left(\sup_{f \in \mathcal{F}_{S, \text{per}}^d(s, R)} \sum_{|\ell|_\infty \leq (m-1)/2} \sum_{k \in \mathbb{Z}^d \setminus \{0\}} |\ell + km|_\infty^{2s} |\langle f, \varphi_{\ell+km} \rangle|^2 \right) \times
\end{aligned}$$

$$\begin{aligned}
& \left(\sup_{|\ell|_\infty \leq (m-1)/2} \sum_{k \in \mathbb{Z}^d \setminus \{0\}} |\ell + km|_\infty^{-2s} \right) \\
&= R^2 m^{-2s} \sup_{|\ell|_\infty \leq (m-1)/2} \sum_{k \in \mathbb{Z}^d \setminus \{0\}} |k + \ell/m|_\infty^{-2s} \\
&\leq R^2 m^{-2s} \left(2^{2s} (2^d - 1) + \sum_{k \in \mathbb{Z}^d \setminus \{0\}} |k|_\infty^{-2s} \right).
\end{aligned}$$

Hence, using Theorem 2.4 we have proved the following result, which extends the scalar results by Brown and Low (1996) and more specifically Rohde (2004) to any dimension $d \geq 1$.

2.8 Theorem. *For d -dimensional periodic Sobolev classes $\mathcal{F}_{S,per}^d(s, R)$ with regularity $s > d/2$ and equidistant design on the cube $[0, 1]^d$ the nonparametric regression experiment $\mathbb{E}_n^d$ and the Gaussian shift experiment $\mathbb{G}_n^d$ are asymptotically equivalent as $n \rightarrow \infty$. The Le Cam distance satisfies*

$$\Delta_{\mathcal{F}_{S,per}^d(s,R)}(\mathbb{E}_n^d, \mathbb{G}_n^d) \lesssim \sigma^{-1} R n^{1/2-s/d}.$$

3 Isomorphic approximation

3.1 General theory

We extend the preceding framework by merely requiring an isomorphic property. Since it will suffice for the subsequent applications, we specialize here immediately to $D_n g = (g(x_1), \dots, g(x_n))$. Let $S_n \subseteq \mathcal{L}^2(\mathcal{D})$, $\dim S_n = n$, have the property

$$\forall g_n \in S_n : g_n(x_1) = \dots = g_n(x_n) = 0 \implies g_n = 0. \quad (3.1)$$

Let

$$\langle f, g \rangle_n := \frac{1}{n} \sum_{i=1}^n f(x_i) \overline{g(x_i)}, \text{ resp. } \langle v, g \rangle_n := \frac{1}{n} \sum_{i=1}^n v_i \overline{g(x_i)}, \quad f, g \in \mathcal{L}^2, v \in \mathbb{R}^n,$$

and $\|g\|_n^2 = \langle g, g \rangle_n$. In this notation Equation (3.1) is equivalent with the isomorphy of the norms $\|\bullet\|_n$ and $\|\bullet\|_{L^2}$ on S_n :

$$\exists A_n, B_n > 0 \forall g_n \in S_n : A_n \|g_n\|_{L^2} \leq \|g_n\|_n \leq B_n \|g_n\|_{L^2}. \quad (3.2)$$

We choose any L^2 -orthonormal basis $(\varphi_j)_{1 \leq j \leq n}$ of S_n and introduce the linear mappings

$$\Pi_n, \mathcal{J}_n : \mathcal{L}^2(\mathcal{D}) \rightarrow S_n, \quad \Pi_n g := \sum_{j=1}^n \langle g, \varphi_j \rangle_n \varphi_j, \quad \mathcal{J}_n g := \Pi_n|_{S_n}^{-1} \Pi_n g.$$

Observe the following properties: for $g_n, h_n \in S_n$ we have $\langle \Pi_n g_n, h_n \rangle = \langle g_n, h_n \rangle_n$ and $\|\Pi_n|_{S_n}\| \leq B_n$, $\|(\Pi_n|_{S_n})^{-1}\| \leq A_n^{-1}$; $\mathcal{J}_n$ is a projection onto S_n and $\mathcal{J}_n g$ interpolates g at the design points (x_i) ; Π_n and $\mathcal{J}_n$ are independent of the choice of the basis (φ_j) .

The regression experiment $\mathbb{E}_n^d$ can be transformed to a functional Gaussian shift by expanding the observations (Y_i) in the basis (φ_j) :

$$Z_1 := \sum_{j=1}^n \langle Y, \varphi_j \rangle_n \varphi_j = \Pi_n f + \frac{\sigma}{\sqrt{n}} (\Pi_n|_{S_n})^{1/2} \zeta \in S_n, \quad (3.3)$$

with Gaussian white noise $\zeta := (\Pi_n|_{S_n})^{-1/2} (\sqrt{n} \sum_{j=1}^n \langle \varepsilon_j, \varphi_j \rangle_n \varphi_j) \sim \mathcal{N}(0, \text{Id}_{S_n})$ because

$$E[\langle \zeta, g_n \rangle \langle \zeta, h_n \rangle] = \langle (\Pi_n|_{S_n})^{-1/2} g_n, (\Pi_n|_{S_n})^{-1/2} h_n \rangle_n = \langle g_n, h_n \rangle_n, \quad g_n, h_n \in S_n.$$

By applying $(\Pi_n|_{S_n})^{-1/2}$ and $(\Pi_n|_{S_n})^{-1}$, respectively, we conclude that the regression experiment $\mathbb{E}_n^d$ is also equivalent to observing

$$Z_2 = (\Pi_n|_{S_n})^{-1/2} Z_1 = (\Pi_n|_{S_n})^{1/2} \mathcal{J}_n f + \frac{\sigma}{\sqrt{n}} \zeta \in S_n, \quad (3.4)$$

$$Z_3 = (\Pi_n|_{S_n})^{-1} Z_1 = \mathcal{J}_n f + \frac{\sigma}{\sqrt{n}} (\Pi_n|_{S_n})^{-1/2} \zeta \in S_n \quad (3.5)$$

with $\zeta \sim \mathcal{N}(0, \text{Id}_{S_n})$.

3.1 Theorem. *The regression experiment $\mathbb{E}^d$ is equivalent with each of the experiments given by observing Z_1 in (3.3), Z_2 in (3.4) and Z_3 in (3.5), respectively.*

The Le Cam distance between $\mathbb{E}_n^d$ and $\mathbb{G}_n^d$ for the class $\mathcal{F}^d$ satisfies the bounds

$$\Delta_{\mathcal{F}^d}(\mathbb{E}_n^d, \mathbb{G}_n^d) \leq 1 - 2\Phi\left(-\frac{\sqrt{n}}{2\sigma} \sup_{f \in \mathcal{F}^d} \|f - \Pi_n|_{S_n}^{1/2} \mathcal{J}_n f\|_{L^2}\right), \quad (3.6)$$

$$\Delta_{\mathcal{F}^d}(\mathbb{E}_n^d, \mathbb{G}_n^d) \leq 1 - 2\Phi\left(-\frac{\sqrt{n}}{2\sigma} \sup_{f \in \mathcal{F}^d} \|f - \mathcal{J}_n f\|_{L^2}\right) + \sqrt{2}\|(\Pi_n|_{S_n})^{-1} - \text{Id}_{S_n}\|_{HS}, \quad (3.7)$$

where $\|\bullet\|_{HS}$ denotes the Hilbert-Schmidt norm of an operator.

Proof. It remains to prove the second part. The first bound (3.6) follows from the equivalence with observing Z_2 by the same arguments as for Theorem 2.4. To establish (3.7), we use the fact that the Hellinger distance between two multivariate normal distributions with the same mean satisfies

$$H^2(N(\mu, \alpha\Sigma), N(\mu, \alpha \text{Id}_{\mathbb{R}^n})) \leq 2\|\Sigma - \text{Id}_{\mathbb{R}^n}\|_{HS}^2, \quad \Sigma \in \mathbb{R}^{n \times n}, \alpha > 0, \quad (3.8)$$

which follows e.g. from (Brown, Cai, Low, and Zhang 2002, Lemma 3) via the diagonalisation $\Sigma = O^\top \text{diag}(\lambda_1, \dots, \lambda_n)O$ and the property $\|\Sigma - \text{Id}_{\mathbb{R}^n}\|_{HS}^2 = \|O(\Sigma - \text{Id}_{\mathbb{R}^n})O^\top\|_{HS}^2 = \sum_{i=1}^n \lambda_i^2$. Therefore the total variation distance between the laws of Z_3 and $Z_4 := \mathcal{J}_n f + \frac{\sigma}{\sqrt{n}}\zeta$ is bounded by

$$\|\mathcal{L}(Z_3) - \mathcal{L}(Z_4)\|_{TV} \leq H(\mathcal{L}(Z_3), \mathcal{L}(Z_4)) \leq \sqrt{2}\|(\Pi_n|_{S_n})^{-1} - \text{Id}_{S_n}\|_{HS}.$$

The by now standard arguments yield with obvious notation

$$\begin{aligned} \Delta_{\mathcal{F}^d}(\mathbb{E}_n^d, \mathbb{G}_n^d) &= \Delta_{\mathcal{F}^d}(Z_3, \mathbb{G}_n^d) \leq \Delta_{\mathcal{F}^d}(Z_4, \mathbb{G}_n^d) + \Delta_{\mathcal{F}^d}(Z_4, Z_3) \\ &\leq 1 - 2\Phi\left(-\frac{\sqrt{n}}{2\sigma} \sup_{f \in \mathcal{F}^d} \|f - \mathcal{J}_n f\|_{L^2}\right) + \sqrt{2}\|(\Pi_n|_{S_n})^{-1} - \text{Id}_{S_n}\|_{HS}, \end{aligned}$$

as asserted. $\square$

3.2 Linear spline approximation

Let us briefly expose how the approach by Carter (2006) fits into the isomorphic framework. As in Section 2.3, we consider equidistant design points $(k/m)_{k \in \{1, \dots, m\}^d}$ with $m = n^{1/d} \in \mathbb{N}$ and periodic functions on the unit cube

$\mathcal{D} = [0, 1]^d$. The space S_n is spanned by the periodized and tensorized linear B-splines

$$b_k(x) = b_k(x_1, \dots, x_d) = \prod_{r=1}^d \bar{b}(mx_r - k_r \bmod 1), \quad \bar{b} := \mathbf{1}_{[-1/2, 1/2]} * \mathbf{1}_{[-1/2, 1/2]},$$

indexed by $k \in \{1, \dots, m\}^d$. For $\alpha \in (1, 2]$ it is well known (cf. De Boor (2001)) that interpolation on S_n for the periodic Hölder class

$$\mathcal{F}_{H, per}^d(\alpha, R) := \{f \in C^\alpha(\mathbb{R}^d) \mid f \text{ } \mathbb{Z}^d\text{-periodic, } \sup_{x \neq y} |\nabla f(x) - \nabla f(y)| / |x - y|^{\alpha-1} \leq R\}$$

satisfies the estimate

$$\sup_{f \in \mathcal{F}_{H, per}^d(\alpha, R)} \|f - \mathcal{I}_n f\|_{L^2([0, 1]^d)} \lesssim R n^{-\alpha/d}. \quad (3.9)$$

On the other hand, we have for $g_n \in S_n$

$$\|g_n\|_{L^2}^2 = \left\| \sum_{k \in \{1, \dots, m\}^d} g_n(k/m) b_k \right\|_{L^2}^2 = \sum_{k, \ell \in \{1, \dots, m\}^d} \langle b_k, b_\ell \rangle_{L^2} g_n(k/m) g_n(\ell/m)$$

with $\langle b_k, b_\ell \rangle_{L^2} = 0$ for $|k - \ell|_\infty > 1$ and $\langle b_k, b_\ell \rangle_{L^2} = 4^{\#\{r: k_r = \ell_r\}} / (6^d n)$ for $|k - \ell|_\infty \leq 1$. Since $\sum_\ell \langle b_k, b_\ell \rangle_{L^2} = \langle b_k, 1 \rangle_{L^2} = n^{-1}$, a weighted Cauchy-Schwarz inequality yields

$$\|g_n\|_{L^2}^2 \leq n^{-1} \sum_{k \in \{1, \dots, m\}^d} g_n(k/m)^2 = \langle g_n, g_n \rangle_n = \langle \Pi_n g_n, g_n \rangle_{L^2}$$

and we conclude, using the ordering of symmetric operators, that $(\Pi_n|_{S_n})^{-1} \leq \text{Id}_{S_n}$.

Adding to the observation Z_3 in (3.5) independent Gaussian noise $\eta \sim \mathcal{N}(0, \frac{\sigma^2}{n}(\text{Id}_{S_n} - (\Pi_n|_{S_n})^{-1}))$, we infer that the regression experiment $\mathbb{E}_n^d$ is more informative than observing

$$Z_5 := Z_3 + \eta = \mathcal{I}_n f + \frac{\sigma}{\sqrt{n}} \tilde{\zeta} \in S_n \quad (3.10)$$

with Gaussian white noise $\tilde{\zeta} := (\Pi_n|_{S_n})^{-1/2} \zeta + n^{1/2} \sigma^{-1} \eta \sim \mathcal{N}(0, \text{Id}_{S_n})$. This randomization together with estimate (3.9) shows that the regression experiment $\mathbb{E}_n^d$ is asymptotically at least as informative as the Gaussian experiment $\mathbb{G}_n^d$ on Hölder

classes $\mathcal{F}_{per}^d(\alpha, R)$ with $\alpha > d/2$ and $d \in \{1, 2, 3\}$. Together with an (easier) randomization in the other direction and a more sophisticated boundary treatment for non-periodic function classes this reproduces the proof by Carter (2006) for asymptotic equivalence of regression and white noise experiments in dimensions 2 and 3. For B-splines of higher order the interpolation property $b_k(i/m) = \delta_{k,i}$ gets lost and $(\Pi_n|_{S_n})^{-1} \leq \text{Id}_{S_n}$ cannot be shown such that a more refined analysis is needed. This will be accomplished in the next section for a similar approach using compactly supported wavelets.

3.3 Wavelet multiresolution analysis

The construction. Let us assume an equidistant dyadic design $(k2^{-j})_{k \in \{1, \dots, 2^j\}^d}$ with $n = 2^{dj}$ points for some $j \in \mathbb{N}$ and $\mathcal{D} = [0, 1]^d$. We consider a wavelet multiresolution analysis $(V_j)_{j \geq 0}$ on $L^2([0, 1]^d)$ obtained by periodisation and tensor products. Let $\bar{\varphi}$ be a standard orthonormal scaling function of an r -regular multiresolution analysis for $L^2(\mathbb{R})$, that is $(\bar{\varphi}(\bullet + k))_{k \in \mathbb{Z}}$ forms an orthonormal system in $L^2(\mathbb{R})$ and satisfies $\int \bar{\varphi} = 1$ as well as the polynomial exactness condition that $\sum_{k \in \mathbb{Z}} k^q \bar{\varphi}(x - k) - x^q$ is a polynomial of maximal degree $q - 1$ for all $q = 0, \dots, R - 1$ (Cohen 2000, Thm. 16.1). We suppose that $\bar{\varphi}$ has compact support in $[-S + 1, S]$, like in Daubechies's construction, so that the functions $\varphi_{jk} : [0, 1]^d \rightarrow \mathbb{R}$, $j \geq 1$, $k \in \{1, \dots, 2^j\}^d$, with

$$\varphi_{jk}(x_1, \dots, x_d) := \sum_{m \in \mathbb{Z}^d} 2^{jd/2} \prod_{i=1}^d \bar{\varphi}(2^j x_i - k_i + 2^j m_i)$$

are well defined and form an orthonormal system in $L^2([0, 1]^d)$ (Wojtaszczyk 1997, Prop. 2.21). We set $S_{2^{jd}} := V_j := \text{span}\{\varphi_{jk} \mid k \in \{1, \dots, 2^j\}^d\}$.

Periodic approximation. Polynomial exactness and continuity of $\bar{\varphi}$ imply for $q = 0, \dots, R - 1$ and any $x \in \mathbb{R}$ (Sweldens and Piessens 1993)

$$\sum_{m \in \mathbb{Z}} (x + m)^q \bar{\varphi}(x + m) = \int_{-\infty}^{\infty} x^q \bar{\varphi}(x) dx.$$

This identity is fundamental for our purposes because it implies for $\mathbb{Z}^d$ -periodic functions $h : \mathbb{R}^d \rightarrow \mathbb{R}$ that coincide with a polynomial p of maximal degree $R - 1$ on $\prod_{i=1}^d [2^{-j}(k_i - S - 1), 2^{-j}(k_i + S)]$:

$$\begin{aligned}
\langle h, \varphi_{jk} \rangle_{L^2} &= \sum_{m \in \mathbb{Z}^d} 2^{jd/2} \int_{[0,1]^d} h(x) \prod_{i=1}^d \bar{\varphi}(2^j(x_i + m_i) - k_i) dx \\
&= 2^{jd/2} \int_{\mathbb{R}^d} h(x) \prod_{i=1}^d \bar{\varphi}(2^j x_i - k_i) dx \\
&= 2^{-jd/2} \int_{[-S-1, S]^d} p(2^{-j}(x + k)) \prod_{i=1}^d \bar{\varphi}(x_i) dx \\
&= 2^{-jd/2} \sum_{m \in \mathbb{Z}^d} p(2^{-j}(m + k)) \prod_{i=1}^d \bar{\varphi}(m_i) \\
&= 2^{-jd/2} \sum_{m \in \{1, \dots, 2^j\}^d} h(2^{-j}m) \varphi_{jk}(2^{-j}m) \\
&= n^{1/2} \langle h, \varphi_{jk} \rangle_n,
\end{aligned}$$

where we identified $n = 2^{jd}$. For any $\mathbb{Z}^d$ -periodic function $g \in H_{S,per}^s([0,1]^d)$ with $s \in (d/2, R)$ this local polynomial reproduction property implies by standard, but sophisticated arguments for direct estimates (Cohen 2000, Thm. 30.6)

$$\|g - \Pi_n g\|_{L^2} \lesssim 2^{-js} \|g\|_{H^s} = n^{-s/d} \|g\|_{H^s}, \quad (3.11)$$

where $\|\bullet\|_{H^s}$ denotes the standard L^2 -Sobolev norm of regularity s on $[0,1]^d$. We split the bias term and obtain by functional calculus

$$\begin{aligned}
\|f - \Pi_n|_{S_n}^{-1/2} \Pi_n f\|_{L^2} &\leq \|f - \Pi_n f\|_{L^2} + \|\Pi_n f - \Pi_n|_{S_n}^{-1/2} \Pi_n f\|_{L^2} \\
&= \|f - \Pi_n f\|_{L^2} + \|h(\Pi_n|_{S_n})(\text{Id} - \Pi_n) \Pi_n f\|_{L^2}
\end{aligned}$$

with $h : \mathbb{R}^+ \rightarrow \mathbb{R}$, $h(x) := 1/(x + x^{1/2}) = (x^{-1/2} - 1)/(1 - x)$. Since h decreases monotonically and $h(x) \leq x^{-1/2}$, we have $\|h(\Pi_n|_{S_n})\|_{L^2 \rightarrow L^2} \leq \lambda_{min}^{-1/2}$ with the smallest eigenvalue λ_{min} of $\Pi_n|_{S_n}$.

$\Pi_n|_{S_n}$ satisfies for $n = 2^{jd} \geq 2S - 1$ the following scaling property:

$$\begin{aligned} \langle \Pi_n \varphi_{jk}, \varphi_{j\ell} \rangle_{L^2} &= \frac{1}{n} \sum_{\nu \in \{1, \dots, 2^j\}^d} \varphi_{jk}(\nu 2^{-j}) \varphi_{j\ell}(\nu 2^{-j}) \\ &= \sum_{m \in \mathbb{Z}^d} \sum_{\nu \in \{1, \dots, 2^j\}^d} \prod_{a=1}^d \left(\bar{\varphi}((\nu - k + 2^j m)_a) \bar{\varphi}((\nu - \ell + 2^j m)_a) \right) \\ &= \prod_{a=1}^d \left(\sum_{b \in \mathbb{Z}} \bar{\varphi}(b - k_a) \bar{\varphi}(b - \ell_a) \right). \end{aligned}$$

Since $\bar{\varphi}$ has compact support, the series is just a finite sum and Π_n has a bounded Toeplitz matrix representation in terms of (φ_{jk}) . Using Fourier multipliers it follows that $\langle \Pi_n g_n, g_n \rangle_{L^2} \geq A_{\bar{\varphi}}^2 \|g_n\|_{L^2}^2$, $g_n \in S_n$, with $A_{\bar{\varphi}} := \inf_{u \in [0, 2\pi]} |\sum_{k \in \mathbb{Z}} \bar{\varphi}(k) e^{iku}|^d$, independently of n . Due to the compact support of $\bar{\varphi}$, we have $A_{\bar{\varphi}} > 0$ iff the trigonometric polynomial $\sum_{k \in \mathbb{Z}} \bar{\varphi}(k) e^{iku}$, $u \in [0, 2\pi]$, does not vanish. It is well known (Sweldens and Piessens 1993, Lemma 3) that this is exactly the condition to ensure that the multiresolution analysis is also generated by an interpolating scaling function. It can be checked for standard Daubechies scaling functions, e.g. by showing $|\bar{\varphi}(k_0)| > \sum_{k' \neq k_0} |\bar{\varphi}(k')|$ for some $k_0 \in \mathbb{Z}$. Moreover, gaining more flexibility by considering the shifted spaces based on $\bar{\varphi}_\tau = \bar{\varphi}(\bullet - \tau)$, $\tau \in (0, 1)$, a wavelet multiresolution analysis will almost always satisfy $A_{\bar{\varphi}_\tau} > 0$ for some value of τ , cf. Sweldens and Piessens (1993) and the references therein.

We arrive at

$$\|f - \Pi_n|_{S_n}^{-1/2} \Pi_n f\|_{L^2} \leq \|f - \Pi_n f\|_{L^2} + A_{\bar{\varphi}}^{-1/2} \|(\text{Id} - \Pi_n) \Pi_n f\|_{L^2}.$$

Because of $\|\Pi_n f\|_{H^s} \rightarrow \|f\|_{H^s}$ (Cohen 2000, Thm. 30.7) we derive from (3.11) the uniform estimate over $f \in \mathcal{F}_{S, \text{per}}^d(s, R)$

$$\|f - \Pi_n|_{S_n}^{-1/2} \Pi_n f\|_{L^2} \leq \|f - \Pi_n f\|_{L^2} + A_{\bar{\varphi}}^{-1/2} \|(\text{Id} - \Pi_n) \Pi_n f\|_{L^2} \lesssim R n^{-s/d}.$$

Hence, the estimate in (3.6) yields asymptotic equivalence between the regression and the white noise experiment for any class $\mathcal{F}_{S, \text{per}}^d(s, R)$ with $s > d/2$.

This result provides another way for constructing explicitly the transformation between the regression and the white noise setting. It has no more theoretical

implications than the Fourier basis approach, but it paves the way for proving asymptotic equivalence for non-periodic function classes.

Non-periodic approximation. Since every φ_{jk} has support length $2^{-j}(2S-1)$, only those functions φ_{jk} with $k_r \in \{1, \dots, S-2\} \cup \{2^j - S + 1, \dots, 2^j\}$ for some $r = 1, \dots, d$ cross the boundary and are periodized at all. Therefore, the same derivation using only interior scaling functions shows that the regression experiment $\mathbb{E}_n^d$ for the general Sobolev function class

$$\mathcal{F}_S^d(s, R) := \{f \in H^s([0, 1]^d) \mid \|f\|_{H^s} \leq R\}$$

is asymptotically more informative than the restricted white noise experiment $\bar{\mathbb{G}}_n^d$ given by observing

$$dY(x) = f(x) dx + \frac{\sigma}{\sqrt{n}} dB(x), \quad x \in [\delta_n, 1 - \delta_n]^d \text{ with } \delta_n := (2S-1)n^{-1/d}. \quad (3.12)$$

Although $\bar{\mathbb{G}}_n^d$ is a priori less informative than $\mathbb{G}_n^d$, we may use classical extrapolation, e.g. the Taylor polynomial T_f^y of order $\lfloor s \rfloor$ around $y \in [\delta_n, 1 - \delta_n]^d$. We define at the points $x \in [0, 1]^d \setminus [\delta_n, 1 - \delta_n]^d$ the extrapolation $\tilde{f}(x) = T_f^{y_x}(x)$ for a point $y_x \in [\delta_n, 1 - \delta_n]^d$ with $|y_x - x|_\infty \leq 2\delta_n$, selected in a measurable way, and $\tilde{f}(x) = f(x)$ otherwise. We thereby achieve

$$\left(\int_{[0, 1]^d} |\tilde{f}(x) - f(x)|^2 dx \right)^{1/2} \lesssim R n^{-s/d}$$

such that

$$\Delta_{\mathcal{F}_S^d(s, R)}(\bar{\mathbb{G}}_n^d, \mathbb{G}_n^d) \lesssim \sigma^{-1} R n^{1/2-s/d}.$$

This means that $\bar{\mathbb{G}}_n^d$ and $\mathbb{G}_n^d$ are asymptotically equivalent for $s > d/2$ and we have obtained a result for function classes without periodicity condition.

3.2 Theorem. *For general d -dimensional Sobolev classes $\mathcal{F}_S^d(s, R)$ with regularity $s > d/2$ and equidistant design on the cube $[0, 1]^d$ the nonparametric regression experiment $\mathbb{E}_n^d$ and the Gaussian white noise experiment $\mathbb{G}_n^d$ are asymptotically*

equivalent as $n \rightarrow \infty$. The Le Cam distance satisfies

$$\Delta_{\mathcal{F}_S^d(s,R)}(\mathbb{E}_n^d, \mathbb{G}_n^d) \lesssim \sigma^{-1} R n^{1/2-s/d}.$$

Discussion. The property that a wavelet estimator based on an equidistant regression model and a corresponding estimator based on a white noise model are asymptotically close is well known, see e.g. Donoho and Johnstone (1999) and Johnstone and Silverman (2004). Interestingly, both papers show identical asymptotics of the L^2 -risk for standard estimators uniformly over balls in Besov spaces $B_{p,q}^s([0,1])$ with $s > 1/p$ or $s = p = 1$. Since $B_{p,q}^s$ embeds into the Sobolev space H^σ for $s > \sigma$ and $s - 1/p > \sigma - 1/2$, Theorem 3.2 provides more generally asymptotic equivalence for Besov classes with $s > 1/p$ and $p < 2$. The counterexample in Brown and Low (1996) shows, however, that for $s \leq 1/2$ and all $p \in [1, \infty]$, asymptotic equivalence breaks down. Similarly, if $\psi \in B_{1,1}^1$ is a function with support in $(0,1)$ and $\|\psi\|_{L^2} = 1$, then $\psi_n(x) := \psi(nx)$ has support in $(0,1/n)$, L^2 -norm $\|\psi_n\|_{L^2} = n^{-1/2}$ and Besov norm $\|\psi_n\|_{B_{1,1}^1} \sim 1$. Hence, testing the signal $f = 0$ versus $f = \psi_n$ has nontrivial power in the white noise model $\mathbb{G}_n^1$, while both signals generate exactly the same observations in the regression model $\mathbb{E}_n^1$. We conclude that $\mathbb{G}_n^1$ and $\mathbb{E}_n^1$ are not asymptotically equivalent on Besov classes with $s = 1$, $p = 1$. An intriguing example for the important class of bounded variation functions is given by $\psi_n(x) = \sqrt{2}\mathbf{1}_{[1/4n, 3/4n]}(x)$. Asymptotic equivalence between Gaussian regression and white noise is indeed an L^2 -theory and we cannot gain by measuring smoothness in an L^p -sense, $p \neq 2$.

Let us also mention that the (asymptotically negligible) loss in information due to neglecting boundary coefficients in the construction seems unavoidable. The wavelets on an interval (Cohen, Daubechies, and Vial 1993) use nonorthogonal boundary corrections and can therefore not be used, while the coiflet approach by Johnstone and Silverman (2004) also involves some information loss at the boundary, cf. their remark on dimensions before Proposition 2.

4 Random design

4.1 The general idea

Denote by $U([0, 1]^d)$ the uniform distribution on the cube $\mathcal{D} = [0, 1]^d$.

4.1 Definition. Let $\mathbb{E}_{n,r}^d$ be the compound experiment obtained from observing independent random design points $X_i \sim U([0, 1]^d)$, $i = 1, \dots, n$, and the regression

$$Y_i = f(X_i) + \sigma \varepsilon_i, \quad i = 1, \dots, n,$$

for $n \in \mathbb{N}$ and $f : [0, 1]^d \rightarrow \mathbb{R}$ in some class $\mathcal{F}^d \subseteq \mathcal{L}^2([0, 1]^d)$ and with i.i.d. random variables $\varepsilon_i \sim \mathcal{N}(0, 1)$, independent of the design.

We place ourselves into the isomorphic setting, that is we are given an $L^2([0, 1]^d)$ -orthonormal basis $(\varphi_j)_{j \geq 1}$ and we set $S_n = \text{span}(\varphi_1, \dots, \varphi_n)$. For the moment we merely assume that S_n is chosen to satisfy the isomorphic condition (3.1) given the random design points $(X_i)_{1 \leq i \leq n}$. Later, certain parts will rely on fine properties of the Fourier basis. Conditionally on the design the regression experiment is equivalent to observing

$$Z_1 := \sum_{j=1}^n \langle Y, \varphi_j \rangle_n \varphi_j = \Pi_n f + \frac{\sigma}{\sqrt{n}} (\Pi_n|_{S_n})^{1/2} \zeta \in S_n$$

with white noise $\zeta \sim N(0, \text{Id}_{S_n})$. Let us briefly comment why the foregoing approaches using Z_2 in (3.4) or Z_3 in (3.5) will not succeed here. For $Z_2 = (\Pi_n|_{S_n})^{-1/2} Z_1$ we need to have $\|(\Pi_n|_{S_n}^{1/2} - \text{Id}) \mathcal{J}_n f\|_{L^2}$ and $\|\mathcal{J}_n f - f\|_{L^2}$ of smaller order than $n^{-1/2}$. The second property can be ensured for Sobolev classes of regularity $s > d/2$ as before. The first property, however, will not hold. By empirical process theory, we have for $g_1, g_2 \in S_n$ approximately $\langle \Pi_n g_1, g_2 \rangle_{L^2} = \langle g_1, g_2 \rangle_n \approx \langle g_1, g_2 \rangle_{L^2} + n^{-1/2} \int g_1 g_2 dB^0$ with a Brownian bridge B^0 . By the linearisation $(1+h)^{1/2} - 1 \approx h/2$ and taking expectation with respect to the random design, we

find

$$\begin{aligned} E\left[\|((\Pi_n|_{S_n})^{1/2} - \text{Id})\mathcal{J}_n f\|_{L^2}^2\right] &\sim E\left[\sum_{j=1}^n \left|n^{-1/2} \int (\mathcal{J}_n f) \varphi_j dB^0\right|^2\right] \\ &\sim n^{-1} \sum_{j=1}^n \int |\varphi_j|^2 |\mathcal{J}_n f|^2 \gtrsim \|\mathcal{J}_n f\|_{L^2}^2. \end{aligned}$$

Hence, in the mean over the random design this term does not tend to zero. When considering $Z_3 = (\Pi_n|_{S_n})^{-1} Z_1$, we would need $\|(\Pi_n|_{S_n})^{-1} - \text{Id}_{S_n}\|_{HS} \rightarrow 0$, compare Bound (3.7), but the mean over this term is by the same approximations of order n . The main defect in these approaches is that we do not take advantage of the regularity of f .

The new idea is based on a two-level procedure, which can be interpreted as a localisation approach, cf. Nussbaum (1996). We choose an intermediate level $n_0 < n$ and split $S_n = S_{n_0} + U_{n_0}^n$ with the $\|\bullet\|_n$ -orthogonal complement $U_{n_0}^n$ of S_{n_0} in S_n . On the low-frequency space S_{n_0} we use the empirical orthogonal projection $P_{n_0}^n Y$ of the data onto S_{n_0} . This construction is analogous to Z_3 in (3.5) and the heteroskedasticity in the noise term will become asymptotically negligible provided $n_0 = o(n^{1/2})$.

On the high-frequency part $U_{n_0}^n$ of S_n we transform to a Gaussian shift with white noise, which is independent of the noise in S_{n_0} , in the spirit of Z_2 in (3.4). In order to take advantage of the regularity of f , however, we do not use the standard square root operator $\Pi_n^{-1/2}$ to whiten the noise, but the adjoint T^* of an operator $T : S_n \rightarrow S_n$ which has an upper triangular matrix representation in the basis (φ_j) and satisfies $TT^* = (\Pi_n|_{S_n})^{-1}$ (as in the Cholesky decomposition). Since T^* is a unitary transformation of $(\Pi_n|_{S_n})^{-1/2}$, the noise part remains white. Due to the triangular structure, the signal coefficients $\langle T^* \Pi_n f, \varphi_j \rangle_{L^2} = \langle T^{-1} \mathcal{J}_n f, \varphi_j \rangle_{L^2}$ do not involve the (usually large) coefficients $\langle \mathcal{J}_n f, \varphi_k \rangle_{L^2}$ for indices k smaller than j . Moreover, for the Fourier basis the other off-diagonal matrix entries of T^{-1} are centred and uncorrelated, while the deviations in the diagonal entries grow with the frequencies, but are exactly counter-balanced by the decay of the Fourier coefficients

for Sobolev function classes. Provided $n_0 \rightarrow \infty$, this high-frequency transformation will imply asymptotic equivalence.

4.2 The main result

Let us specify the transformation T concretely based on the Gram-Schmidt procedure for orthonormalisation with respect to $\|\bullet\|_n$. For $j \leq n$ denote by $P_j, P_j^n : S_n \rightarrow S_n$ the L^2 -orthogonal and $\|\bullet\|_n$ -orthogonal projections onto S_j , respectively, and set $P_0^n := 0$. We obtain an $\|\bullet\|_n$ -orthonormal basis (φ_j^n) of S_n via

$$\varphi_j^n := \frac{\varphi_j - P_{j-1}^n \varphi_j}{\|\varphi_j - P_{j-1}^n \varphi_j\|_n}, \quad j = 1, \dots, n.$$

Then φ_j^n is in S_j and the $\|\bullet\|_n$ -orthogonality $\varphi_j^n \perp_n S_{j-1}$ holds. Defining $T : S_n \rightarrow S_n$ via $T\varphi_j := \varphi_j^n$, we see that T satisfies $\langle T\varphi_{j'}, \varphi_j \rangle_{L^2} = 0$ for $j > j'$ and is an isometry between $(S_n, \|\bullet\|_{L^2})$ and $(S_n, \|\bullet\|_n)$ such that $\Pi_n|_{S_n} = (TT^*)^{-1}$. The noise terms $(\langle \varepsilon, \varphi_j^n \rangle_n)_{1 \leq j \leq n} \sim \mathcal{N}(0, n^{-1})$ are therefore independent and

$$P_{n_0}^n \varepsilon := \sum_{j=1}^{n_0} \langle \varepsilon, \varphi_j^n \rangle_n \varphi_j^n = \sum_{j=1}^{n_0} \langle \varepsilon, \varphi_j^n \rangle_n T\varphi_j \sim \mathcal{N}(0, n^{-1} T|_{S_{n_0}} T|_{S_{n_0}}^*).$$

Using $T|_{S_{n_0}} T|_{S_{n_0}}^* = (\Pi_n|_{S_{n_0}})^{-1}$, we introduce the rescaled covariance operator $\Sigma : S_n \rightarrow S_n$ via

$$\Sigma g_n := (\Pi_n|_{S_{n_0}})^{-1} P_{n_0} g_n + (\text{Id}_{S_n} - P_{n_0}) g_n, \quad g_n \in S_n.$$

The regression experiment is then transformed to observing

$$\begin{aligned} Z_r &:= \sum_{j=1}^{n_0} \langle Y, \varphi_j^n \rangle_n \varphi_j^n + \sum_{j=n_0+1}^n \langle Y, \varphi_j^n \rangle_n \varphi_j \in S_n \\ &= P_{n_0}^n f + T^{-1}(P_n^n - P_{n_0}^n) f + n^{-1/2} \sigma \Sigma^{1/2} \zeta \in S_n \end{aligned} \quad (4.1)$$

with Gaussian white noise $\zeta \sim N(0, \text{Id}_{S_n})$, conditional on the random design.

4.2 Example. Let us consider the Haar basis. Write $I_{jk} = [2^{-j}k, 2^{-j}(k+1))$, $N_{jk} = \#\{i : X_i \in I_{jk}\}$ and $\psi_{jk} = 2^{j/2}(\mathbf{1}_{I_{j+1,2k}} - \mathbf{1}_{I_{j+1,2k+1}})$ for $j \geq 0, k =$

$0, \dots, 2^j - 1$. By construction the transformed basis function ψ_{jk}^n has support I_{jk} , is constant on $I_{j+1,2k}$, $I_{j+1,2k+1}$ and satisfies $\langle \psi_{jk}^n, \mathbf{1}_{I_{jk}} \rangle_n = 0$, $\|\psi_{jk}^n\|_n = 1$. We infer

$$\psi_{jk}^n = C_{jk} (N_{j,2k}^{-1} \mathbf{1}_{I_{j,2k}} - N_{j,2k+1}^{-1} \mathbf{1}_{I_{j,2k+1}}), \quad C_{jk}^2 = n N_{j+1,2k} N_{j+1,2k+1} / N_{jk}.$$

This is exactly the application of our framework underlying previous one-dimensional constructions (Brown, Cai, Low, and Zhang 2002, Eq. (2.8)). Because here S_n is for most design realisations not isomorphic, additional randomisations are needed.

For the following general d -dimensional theorem we consider the construction (4.1) in terms of the Fourier basis functions $\varphi_j(x) = \exp(2\pi i \langle \ell(j), x \rangle)$ with an enumeration $\ell : \mathbb{N} \rightarrow \mathbb{Z}^d$ of $\mathbb{Z}^d$ satisfying $|\ell(j)|_{\ell^2} \leq |\ell(j')|_{\ell^2}$ for $j \leq j'$ (i.e. sorted in the order of magnitudes of the frequencies).

4.3 Theorem. *For d -dimensional periodic Sobolev classes $\mathcal{F}_{S,per}^d(s, R)$ with regularity $s > d/2$ the nonparametric regression experiment $\mathbb{E}_{n,r}^d$ with random design and the Gaussian shift experiment $\mathbb{G}_n^d$ are asymptotically equivalent as $n_0, n \rightarrow \infty$ and $n_0 = o(n^{1/2})$. The Le Cam distance satisfies*

$$\Delta_{\mathcal{F}_{S,per}^d(s,R)}(\mathbb{E}_{n,r}^d, \mathbb{G}_n^d) \lesssim n^{-1/2} n_0 + \sigma^{-1} R n_0^{1/2-s/d}.$$

4.4 Remark. *The asymptotically optimal choice of n_0 is given by $n_0 \sim n^{d/(2s+d)}$, which yields a bound on the Le Cam distance of order $n^{(d-2s)/(2d+4s)}$. Note that this choice $n_0 \sim n^{d/(2s+d)}$ corresponds exactly to the optimal dimension of the approximation spaces in nonparametric regression and is also used by Gaïffas (2005) for his two-level construction of optimal confidence bands.*

Proof. In order to bound the Le Cam distance for compound experiments, we use that for distributions $K \otimes P$ and $K' \otimes P$, defined on $(\Omega \times \Omega', \mathcal{F} \otimes \mathcal{F}')$ by the measure P on $\mathcal{F}$ and the Markov kernels K, K' from Ω to $\mathcal{F}'$, the total variation distance can be calculated by conditioning:

$$\|K \otimes P - K' \otimes P\|_{TV(\mathcal{F} \otimes \mathcal{F}')} = \int \|K(\omega, \bullet) - K'(\omega, \bullet)\|_{TV(\mathcal{F}')} P(d\omega).$$

Therefore we can first work conditionally on the design and then take expectations for (X_i) . Moreover, the white noise experiment $\mathbb{G}_n^d$ is equivalent to the compound experiment of $\mathbb{G}_n^d$ and the observation of the random design points because the latter is a trivial randomisation of $\mathbb{G}_n^d$.

It is a remarkable property of the Fourier basis that S_n is almost surely isomorphic, cf. Theorem 1.1 in Bass and Gröchenig (2004). In Proposition 4.8 below we prove that the event

$$\Omega_j^n := \{\forall g \in S_j : \frac{1}{2}\|g\|_{L^2} \leq \|g\|_n \leq 2\|g\|_{L^2}\} \quad (4.2)$$

for $j \log(j) = o(n)$ even satisfies $P((\Omega_j^n)^c) \rightarrow 0$ with a convergence rate faster than any polynomial in n . This is much tighter with respect to the subspace dimension than what can be derived from Bass and Gröchenig (2004). In order to establish asymptotic equivalence, it suffices therefore to estimate the total variation distances on the event $\Omega_{n_0}^n$.

By (4.1), the regression experiment $\mathbb{E}_{n,r}^d$ is equivalent to observing Z_r together with the design. Introducing

$$\bar{Z}_r := P_n f + \sigma n^{-1/2} \zeta \quad \in S_n, \quad (4.3)$$

we shall prove in a moment that (with obvious notation)

$$\Delta_{\mathcal{F}_{S,per}^d(s,R)}(Z_r, \bar{Z}_r) \lesssim n^{-1/2} n_0 + \sigma^{-1} R n_0^{1/2-s/d}, \quad (4.4)$$

but then the assertion follows: Observing $\bar{Z}_r$ is equivalent to observing

$$dY(x) = P_n f(x) + \sigma n^{-1/2} dB(x), \quad x \in [0, 1]^d,$$

which has a total variation distance to the Gaussian shift $\mathbb{G}_n^d$ of order $\sigma^{-1} n^{1/2} \|f - P_n f\|_{L^2} \lesssim \sigma^{-1} n^{1/2-s/d} \|f\|_{H^s}$. Using the triangle inequality for the Le Cam distance between the intermediate experiments, we arrive at the bound for $\Delta_{\mathcal{F}_{S,per}^d(s,R)}(\mathbb{E}_{n,r}^d, \mathbb{G}_n^d)$.

To obtain (4.4), we take expectations over the design and split

$$E[\|\mathcal{L}(Z_r) - \mathcal{L}(Z'_r)\|_{TV}^2 \mathbf{1}_{\Omega_{n_0}^n}] \lesssim I + II + III$$

with the terms

$$\begin{aligned} I &:= n\sigma^{-2}E[\|(P_{n_0}^n - P_{n_0})f\|_{L^2}^2 \mathbf{1}_{\Omega_{n_0}^n}] \text{ (difference in mean on } S_{n_0}), \\ II &:= E[\|(\Pi_n|_{S_{n_0}})^{-1} - \text{Id}_{S_{n_0}}\|_{HS}^2 \mathbf{1}_{\Omega_{n_0}^n}] \text{ (heteroskedasticity on } S_{n_0}), \\ III &:= n\sigma^{-2}E[\|(T^{-1}(P_n^n - P_{n_0}^n) - (P_n - P_{n_0}))f\|_{L^2}^2 \mathbf{1}_{\Omega_{n_0}^n}] \text{ (difference in mean on } S_{n_0}^{\perp L^2}). \end{aligned}$$

Term I. Using the projection properties, we obtain on $\Omega_{n_0}^n$:

$$\|(P_{n_0}^n - P_{n_0})f\|_{L^2}^2 = \|P_{n_0}^n(\text{Id} - P_{n_0})f\|_{L^2}^2 \leq 4\|P_{n_0}^n(\text{Id} - P_{n_0})f\|_n^2.$$

Because of $E[\langle \varphi_k, \varphi_j^n \rangle_n \overline{\langle \varphi_{k'}, \varphi_j^n \rangle_n}] = 0$ for $k \neq k'$, $k, k' > j$ by Proposition 4.5 below, an expansion in the basis (φ_j^n) yields

$$\begin{aligned} E[\|P_{n_0}^n(\text{Id} - P_{n_0})f\|_n^2] &= \sum_{j=1}^{n_0} \sum_{k=n_0+1}^{\infty} |\langle f, \varphi_k \rangle_{L^2}|^2 E[|\langle \varphi_k, \varphi_j^n \rangle_n|^2] \\ &= \sum_{k=n_0+1}^{\infty} |\langle f, \varphi_k \rangle_{L^2}|^2 E[\|P_{n_0}^n \varphi_k\|_n^2]. \end{aligned}$$

Proposition 4.9 below yields $E[\|P_{n_0}^n \varphi_k\|_n^2] \lesssim k/n$ and hence

$$I \lesssim \sigma^{-2} \sum_{k=n_0+1}^{\infty} |\langle f, \varphi_k \rangle_{L^2}|^2 k \lesssim \sigma^{-2} n_0^{1-2s/d} \|f\|_{H^s}^2.$$

Term II. Using $\|(\Pi_n|_{S_{n_0}})^{-1}\|_{L^2 \rightarrow L^2} \leq 4$ on $\Omega_{n_0}^n$, we find:

$$\begin{aligned} E[\|(\Pi_n|_{S_{n_0}})^{-1} - \text{Id}_{S_{n_0}}\|_{HS}^2 \mathbf{1}_{\Omega_{n_0}^n}] &\leq E[\|(\Pi_n|_{S_{n_0}})^{-1}\|_{L^2 \rightarrow L^2} \| \Pi_n|_{S_{n_0}} - \text{Id}_{S_{n_0}} \|_{HS}^2 \mathbf{1}_{\Omega_{n_0}^n}] \\ &\leq 4E[\|\Pi_n|_{S_{n_0}} - \text{Id}_{S_{n_0}}\|_{HS}^2] \\ &= 4 \sum_{j,j'=1}^{n_0} E[|\langle \varphi_j, \varphi_{j'} \rangle_n - \delta_{j,j'}|^2] \\ &\leq 4n^{-1} \sum_{j,j'=1}^{n_0} \int |\varphi_j|^2 |\varphi_{j'}|^2. \end{aligned}$$

For the Fourier basis we obtain $II \leq 4n^{-1}n_0^2$.

Term III. Let us write $f = f_0 + f_1 + f_2$ with $f_0 = P_{n_0}f$, $f_1 = (P_n - P_{n_0})f$,

$f_2 = (\text{Id} - P_n)f$. Then the projection properties imply

$$\begin{aligned}
& E[\|(T^{-1}(P_n^n - P_{n_0}^n) - (P_n - P_{n_0}))f\|_{L^2}^2 \mathbf{1}_{\Omega_{n_0}^n}] \\
&= E[\|T^{-1}f_1 + T^{-1}P_n^n f_2 - T^{-1}P_{n_0}^n(f_1 + f_2) - f_1\|_{L^2}^2 \mathbf{1}_{\Omega_{n_0}^n}] \\
&\leq 3E[\|(T^{-1} - \text{Id})f_1\|_{L^2}^2 + \|(P_n^n - P_{n_0}^n)f_2\|_n^2 + \|P_{n_0}^n f_1\|_n^2 \mathbf{1}_{\Omega_{n_0}^n}] \\
&\leq 3E[\|f_1\|_n^2 + \|f_1\|_{L^2}^2 - 2\text{Re}(\langle T^{-1}f_1, f_1 \rangle_{L^2})] + 3E[\|f_2\|_n^2] + 3E[\|P_{n_0}^n f_1\|_n^2 \mathbf{1}_{\Omega_{n_0}^n}] \\
&= 6E[\text{Re}(\langle f_1 - T^{-1}f_1, f_1 \rangle_{L^2})] + 3\|f_2\|_{L^2}^2 + 3E[\|P_{n_0}^n f_1\|_n^2 \mathbf{1}_{\Omega_{n_0}^n}] \\
&=: III_1 + III_2 + III_3.
\end{aligned}$$

The term III_2 is easily bounded by $\|f_2\|_{L^2}^2 \lesssim n^{-2s/d}\|f\|_{H^s}^2$. As in the estimate for term I, we obtain $III_3 \lesssim n^{-1}n_0^{1-2s/d}\|f\|_{H^s}^2$. For III_1 we use $E[\langle T^{-1}\varphi_j, \varphi_k \rangle_{L^2}] = 0$, $j \neq k$, by Proposition 4.5 below to conclude

$$E[\text{Re}(\langle f_1 - T^{-1}f_1, f_1 \rangle_{L^2})] = \sum_{j=n_0+1}^n |\langle f, \varphi_j \rangle_{L^2}|^2 E[\langle (\text{Id} - T^{-1})\varphi_j, \varphi_j \rangle_{L^2}].$$

Because of $\|\varphi_j\|_n = 1$ for the Fourier basis we find

$$\langle T^{-1}\varphi_j, \varphi_j \rangle_{L^2} = \langle \|\varphi_j - P_{j-1}^n \varphi_j\|_n \varphi_j^n + P_{j-1}^n \varphi_j, \varphi_j^n \rangle_n = \|\varphi_j - P_{j-1}^n \varphi_j\|_n \geq 1 - \|P_{j-1}^n \varphi_j\|_n^2.$$

By Proposition 4.9 below, the bound

$$E[\text{Re}(\langle f_1 - T^{-1}f_1, f_1 \rangle_{L^2})] \leq \sum_{j=n_0+1}^n |\langle f, \varphi_j \rangle_{L^2}|^2 E[\|P_{j-1}^n \varphi_j\|_n^2] \lesssim \sum_{j=n_0+1}^n \frac{j}{n} |\langle f, \varphi_j \rangle_{L^2}|^2$$

follows, which is of order $n^{-1}n_0^{1-2s/d}\|f\|_{H^s}^2$. Putting the estimates together, we have shown

$$III \lesssim \sigma^{-2}(n_0^{1-2s/d}\|f\|_{H^s}^2 + n^{1-2s/d}\|f\|_{H^s}^2 + n_0^{1-2s/d}\|f\|_{H^s}^2) \lesssim \sigma^{-2}n_0^{1-2s/d}\|f\|_{H^s}^2$$

and in sum $I + II + III \lesssim \sigma^{-2}n_0^{1-2s/d}R^2 + n^{-1}n_0^2$ uniformly over $f \in \mathcal{F}_{S,per}^d(s, R)$, which gives the asserted bound (4.4). $\square$

4.3 Technical results

We gather results on fine properties of the Fourier basis (φ_j) and its generated approximation spaces S_n . The setting is as in the proof of Theorem 4.3. For the value of the next proposition notice that $\langle \varphi_{k'}, \varphi_k^n \rangle_n = \langle T^{-1}\varphi_{k'}, \varphi_k \rangle_{L^2}$.

4.5 Proposition. *We have for indices $k'', k' > k \geq 1$, $k'' \neq k'$:*

$$E[\langle \varphi_{k'}, \varphi_k^n \rangle_n] = 0 \quad \text{and} \quad E[\langle \varphi_{k'}, \varphi_k^n \rangle_n \overline{\langle \varphi_{k''}, \varphi_k^n \rangle_n}] = 0.$$

Proof. Since the randomness enters via P_{k-1}^n in a very intricate way, we use a symmetry argument. Specify $X_i := (Y_i + \vartheta) \bmod 1$, $i = 1, \dots, n$, with $Y_i \sim U([0, 1]^d)$, $\vartheta \sim U([0, 1]^d)$ all independent such that $X_i \sim U([0, 1]^d)$ i.i.d. Working conditionally on ϑ , we shall keep track on the dependence on ϑ using brackets. We claim that for $k' > k$ it holds that

$$\langle \varphi_{k'}, \varphi_k^n \rangle_n[\vartheta] = e^{2\pi i \langle \ell(k') - \ell(k), \vartheta \rangle} \langle \varphi_{k'}, \varphi_k^n \rangle_n[0], \quad (4.5)$$

which entails the result due to

$$\int_{[0, 1]^d} e^{2\pi i \langle \ell(k') - \ell(k), \vartheta \rangle} d\vartheta = 0 \quad \text{and} \quad \int_{[0, 1]^d} e^{2\pi i (\langle \ell(k') - \ell(k), \vartheta \rangle - \langle \ell(k'') - \ell(k), \vartheta \rangle)} d\vartheta = 0.$$

For $m \in \mathbb{Z}^d$ put

$$A_m[\vartheta] := \frac{1}{n} \sum_{j=1}^n e^{2\pi i \langle m, X_j[\vartheta] \rangle} = \frac{1}{n} \sum_{j=1}^n e^{2\pi i \langle m, Y_j + \vartheta \rangle} = e^{2\pi i \langle m, \vartheta \rangle} A_m[0].$$

The proof of (4.5) will be performed by induction from $\kappa < k$ to k , considering tupels (κ', κ) , $\kappa' > \kappa$, and (k', k) , $k' > k$. Since $\ell(1) = 0$ and $\varphi_1^n = \varphi_1 = 1$, we have for $k' > 1$ and $k = 1$

$$\langle \varphi_{k'}, \varphi_k^n \rangle_n[\vartheta] = \frac{1}{n} \sum_{j=1}^n e^{2\pi i \langle \ell(k'), Y_j + \vartheta \rangle} = e^{2\pi i \langle \ell(k') - \ell(1), \vartheta \rangle} \langle \varphi_{k'}, \varphi_k^n \rangle_n[0].$$

Writing $c_k := \|\varphi_k - P_{k-1}^n \varphi_k\|_n^{-1}$, the induction hypothesis implies

$$c_k^{-2}[\vartheta] = 1 - \sum_{j=1}^{k-1} |\langle \varphi_k, \varphi_j^n \rangle|^2[\vartheta] = c_k^{-2}[0]$$

and furthermore

$$\begin{aligned} \langle \varphi_{k'}, \varphi_k^n \rangle_n[\vartheta] &= \langle \varphi_{k'}, c_k(\varphi_k - \sum_{r=1}^{k-1} \langle \varphi_k, \varphi_r^n \rangle \varphi_r^n) \rangle[\vartheta] \\ &= c_k \left(\langle \varphi_{k'}, \varphi_k \rangle_n - \sum_{r=1}^{k-1} \langle \varphi_{k'}, \varphi_r^n \rangle_n \overline{\langle \varphi_k, \varphi_r^n \rangle_n} \right)[\vartheta] \\ &= c_k[\vartheta] \left(A_{\ell(k') - \ell(k)}[\vartheta] - \sum_{r=1}^{k-1} e^{2\pi i \langle \ell(k') - \ell(k), \vartheta \rangle} \langle \varphi_{k'}, \varphi_r^n \rangle_n[0] \overline{\langle \varphi_k, \varphi_r^n \rangle_n[0]} \right) \\ &= e^{2\pi i \langle \ell(k') - \ell(k), \vartheta \rangle} \langle \varphi_{k'}, \varphi_k^n \rangle_n[0], \end{aligned}$$

which proves the induction step and gives (4.5). $\square$

4.6 Proposition. *Suppose $g = \sum_{|\ell|_{\ell^2} \leq L} \gamma_\ell e^{2\pi i \langle \ell, \bullet \rangle}$ is a d -dimensional trigonometric polynomial of degree L . Let $\Delta \in (0, L^{-1}]$ with $1/\Delta \in \mathbb{N}$ be given and define the cubes $C_m := \prod_{i=1}^d [(m_i - 1)\Delta, m_i\Delta)$. Then:*

$$\Delta^d \sum_{m \in \{1, \dots, \Delta^{-1}\}^d} \sup_{x_m \in C_m} \left| |g(x_m)|^2 - |g(m\Delta)|^2 \right| \leq \|g\|_{L^2}^2 (e^{2d\Delta L} - 1).$$

Proof. We need multi-indices $\alpha, \beta \in \mathbb{N}_0^d$ with $\alpha! := \alpha_1! \cdots \alpha_d!$, $x^\alpha := x_1^{\alpha_1} \cdots x_d^{\alpha_d}$, $\binom{\alpha}{\beta} := \frac{\alpha!}{\beta!(\alpha-\beta)!}$ and differential operators $D^\alpha := \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_d}}{\partial x_d^{\alpha_d}}$. Since $|g|^2$ is real-analytic, a power series expansion gives for any $x_m \in C_m$:

$$\begin{aligned} ||g|^2(x_m) - |g|^2(m\Delta)| &= \left| \sum_{\alpha \in \mathbb{N}_0^d, \alpha \neq 0} D^\alpha |g|^2(m\Delta) \frac{(x_m - m\Delta)^\alpha}{\alpha!} \right| \\ &\leq \sum_{\alpha \in \mathbb{N}_0^d, \alpha \neq 0} \frac{\Delta^{|\alpha|_{\ell^1}}}{\alpha!} \sum_{\beta \in \mathbb{N}_0^d, \beta \leq \alpha} \binom{\alpha}{\beta} |D^\beta g(m\Delta)| |D^{\alpha-\beta} \bar{g}(m\Delta)|. \end{aligned}$$

Together with g any derivative is again a trigonometric polynomial of degree L and by the isometry (2.3) and Bernstein's inequality, cf. (Meyer 1995, p. 32), we obtain

$$\Delta^d \sum_{m \in \{1, \dots, \Delta^{-1}\}^d} |D^\alpha g|^2(m\Delta) = \|D^\alpha g\|_{L^2}^2 \leq L^{2|\alpha|_{\ell^1}} \|g\|_{L^2}^2.$$

This implies by the Cauchy-Schwarz inequality

$$\begin{aligned} &\Delta^d \sum_{m \in \{1, \dots, \Delta^{-1}\}^d} \sup_{x_m \in C_m} ||g|^2(x_m) - |g|^2(m\Delta)| \\ &\leq \Delta^d \sum_{\alpha \in \mathbb{N}_0^d, \alpha \neq 0} \frac{\Delta^{|\alpha|_{\ell^1}}}{\alpha!} \sum_{\beta \in \mathbb{N}_0^d, \beta \leq \alpha} \binom{\alpha}{\beta} \left(\sum_m |D^\beta g(m\Delta)|^2 \right)^{1/2} \left(\sum_m |D^{\alpha-\beta} \bar{g}(m\Delta)|^2 \right)^{1/2} \\ &\leq \|g\|_{L^2}^2 \sum_{\alpha \in \mathbb{N}_0^d, \alpha \neq 0} \frac{\Delta^{|\alpha|_{\ell^1}}}{\alpha!} \sum_{\beta \in \mathbb{N}_0^d, \beta \leq \alpha} \binom{\alpha}{\beta} L^{|\beta|_{\ell^1}} L^{|\alpha-\beta|_{\ell^1}} \\ &= \|g\|_{L^2}^2 \sum_{\alpha \in \mathbb{N}_0^d, \alpha \neq 0} \frac{(2\Delta L)^{|\alpha|_{\ell^1}}}{\alpha!} \\ &= \|g\|_{L^2}^2 (e^{2d\Delta L} - 1). \end{aligned}$$

$\square$

4.7 Lemma. *Let $Y \in \mathbb{R}^r$ follow the multinomial distribution with parameters n and $p_1 = \dots = p_r = 1/r$. Then for $n \rightarrow \infty$ and $r = r(n)$ with $r \log(r)/n \rightarrow 0$*

$$\forall C > 0 : \limsup_{n \rightarrow \infty} \frac{1}{4} r(n)^{C^2/4-1} P\left(\max_{1 \leq i \leq r(n)} |Y_i - n/r(n)| > C \sqrt{n \log(r(n))/r(n)}\right) \leq 1.$$

Proof. If $X_1, \dots, X_r$ are independently Poisson(n/r)-distributed, then it is well known that the law of $(X_1, \dots, X_r)$ given $\sum_{i=1}^r X_i = n$ is multinomial with parameters n and $p_1 = \dots = p_r = 1/r$. Set $A_{nr} := C \sqrt{n \log(r)/r}$. Since

$$k \mapsto P\left(\max_{1 \leq i \leq r} X_i - n/r > A_{nr} \mid \sum_{i=1}^r X_i = k\right)$$

is obviously increasing in $k \in \mathbb{N}$, we obtain

$$P\left(\max_{1 \leq i \leq r} X_i - n/r > A_{nr} \mid \sum_{i=1}^r X_i = n\right) \leq \frac{P(\max_{1 \leq i \leq r} X_i - n/r > A_{nr})}{P(\sum_{i=1}^r X_i \geq n)}.$$

As $\sum_{i=1}^r X_i$ is Poisson(n)-distributed, $\lim_{n \rightarrow \infty} P(\sum_{i=1}^r X_i \geq n) = 1/2$ holds, whence

$$\limsup_{n \rightarrow \infty} \left(P(\max_{1 \leq i \leq r} Y_i - n/r > A_{nr}) - 2P(\max_{1 \leq i \leq r} X_i - n/r > A_{nr}) \right) \leq 0. \quad (4.6)$$

By the exponential moment estimate $E[e^{a(X_i - n/r)}] = e^{n(e^a - a - 1)/r} \leq e^{3na^2/4r}$ for $a := rA_{nr}/n \rightarrow 0$ and n large, the generalized Markov inequality yields

$$P(\max_{1 \leq i \leq r} X_i - n/r > A_{nr}) \leq rP(X_i - n/r > A_{nr}) \leq re^{3na^2/4r - aA_{nr}} = r^{1-C^2/4}.$$

By use of (4.6) and a completely symmetric argument for $P(\max_{1 \leq i \leq r} (n/r - X_i) > A_{nr})$, the result follows. $\square$

4.8 Proposition. *For $j = j(n)$ such that $j \log(j) = o(n)$ and the event Ω_j^n in (4.2) we have $\lim_{n \rightarrow \infty} n^p P((\Omega_{j(n)}^n)^c) = 0$ for any power $p > 0$.*

Proof. From Proposition 4.6 we derive with $\Delta \leq L := |\ell(j)|_{\ell^2}$, $1/\Delta \in \mathbb{N}$, the cubes

$C_m := \prod_{i=1}^d [(m_i - 1)\Delta, m_i\Delta)$ and the occupations $N_m := \#\{i : X_i \in C_m\}$:

$$\begin{aligned}
& \left| \|g\|_{L^2}^2 - \frac{1}{n} \sum_{i=1}^n |g(X_i)|^2 \right| \\
&= \frac{1}{n} \left| \sum_{m \in \{1, \dots, \Delta^{-1}\}^d} \left(\Delta^d n |g(m\Delta)|^2 - \sum_{i: X_i \in C_m} |g(X_i)|^2 \right) \right| \\
&\leq \frac{1}{n} \sum_{m \in \{1, \dots, \Delta^{-1}\}^d} \left(|\Delta^d n - N_m| |g(m\Delta)|^2 + N_m \sup_{x_m \in C_m} \left| |g(m\Delta)|^2 - |g(x_m)|^2 \right| \right) \\
&\leq \frac{\|g\|_{L^2}^2}{\Delta^d n} \max_{m \in \{1, \dots, \Delta^{-1}\}^d} \left(|\Delta^d n - N_m| + N_m (e^{2d\Delta L} - 1) \right) \\
&\leq \|g\|_{L^2}^2 \left(e^{2d\Delta L} \max_{m \in \{1, \dots, \Delta^{-1}\}^d} |1 - N_m/n\Delta^d| + (e^{2d\Delta L} - 1) \right)
\end{aligned}$$

By Lemma 4.7 $\max_m |1 - N_m/n\Delta^d| \geq C(n\Delta^d)^{-1} \log(1/\Delta)$ has probability tending to zero with any given polynomial rate when choosing C sufficiently large. Since $L^d \log(L) \lesssim j \log(j) = o(n)$, we can choose $\Delta = o(L^{-1})$ such that still $\Delta^{-d} \log^2(1/\Delta) = o(n)$ holds. This gives

$$|\|g\|_{L^2}^2 - \|g\|_n^2| \leq \left(C e^{2d\Delta L} (n\Delta^d)^{-1} \log(1/\Delta) + (e^{2d\Delta L} - 1) \right) \|g\|_{L^2}^2 \leq \frac{3}{4} \|g\|_{L^2}^2$$

for large n with probability larger than $1 - n^{-p}$. $\square$

4.9 Proposition. *For $j \in \mathbb{N}$ with $j \log(j) = o(n)$ we have*

$$E[\|P_{j-1}^n \varphi_j\|_n^2] \lesssim j/n.$$

Proof. By construction, $\|P_{j-1}^n \varphi_j\|_n^2 \leq \|\varphi_j\|_n^2 = 1$ holds so that by Proposition 4.8 it suffices to find the bound for the expectation on the event Ω_j^n .

Setting $A_m := \frac{1}{n} \sum_{k=1}^m \exp(2\pi i \langle m, X_k \rangle)$, $m \in \mathbb{Z}^d$, we use Parseval's identity and $E[|A_m|^2] = 1/n$ for $m \neq 0$ to obtain

$$\begin{aligned}
E[\|P_{j-1}^n \varphi_j\|_n^2 \mathbf{1}_{\Omega_j^n}] &= E \left[\sup_{g \in V_{j-1}} \frac{|\langle \varphi_j, g \rangle_n|^2}{\|g\|_n^2} \mathbf{1}_{\Omega_j^n} \right] \\
&\leq E \left[\sup_{\|(c_r)\|_{\ell^2} = 1} \left| \frac{1}{n} \sum_{k=1}^n \sum_{r=1}^{j-1} \bar{c}_r e^{2\pi i \langle \ell(j) - \ell(r), X_k \rangle} \right|^2 \sup_{g \in V_{j-1}} \frac{\|g\|_{L^2}^2}{\|g\|_n^2} \mathbf{1}_{\Omega_j^n} \right]
\end{aligned}$$

$$\leq 4E \left[\sum_{r=1}^{j-1} |A_{\ell(j)-\ell(r)}|^2 \right] = \frac{4(j-1)}{n}.$$

This gives the result. $\square$

References

- BASS, R. F., AND K. GRÖCHENIG (2004): “Random sampling of multivariate trigonometric polynomials.,” *SIAM J. Math. Anal.*, 36(3), 773–795.
- BROWN, L. D., T. CAI, M. G. LOW, AND C.-H. ZHANG (2002): “Asymptotic equivalence theory for nonparametric regression with random design.,” *Ann. Stat.*, 30(3), 688–707.
- BROWN, L. D., AND M. G. LOW (1996): “Asymptotic equivalence of nonparametric regression and white noise.,” *Ann. Stat.*, 24(6), 2384–2398.
- BROWN, L. D., AND C.-H. ZHANG (1998): “Asymptotic nonequivalence of nonparametric experiments when the smoothness index is $1/2$.,” *Ann. Stat.*, 26(1), 279–287.
- CAI, T., AND L. D. BROWN (1999): “Wavelet estimation for samples with random uniform design.,” *Stat. Probab. Lett.*, 42(3), 313–321.
- CARTER, A. (2006): “A continuous Gaussian process approximation to a nonparametric regression in two dimensions.,” *Bernoulli*, 12(1), 143–156.
- COHEN, A. (2000): “Wavelet methods in numerical analysis.,” in *Handbook of numerical analysis. Vol. 7*, ed. by P. G. Ciarlet. North-Holland/ Elsevier, Amsterdam.
- COHEN, A., I. DAUBECHIES, AND P. VIAL (1993): “Wavelets on the interval and fast wavelet transforms.,” *Appl. Comput. Harmon. Anal.*, 1(1), 54–81.
- DALALYAN, A., AND M. REISS (2006): “Asymptotic statistical equivalence for ergodic diffusions: the multidimensional case.,” *Prob. Th. Rel. Fields*, to appear.

- DE BOOR, C. (2001): *A practical guide to splines. Rev. ed.* Applied Mathematical Sciences. 27. New York, NY: Springer.
- DONOHO, D. L., AND I. M. JOHNSTONE (1999): “Asymptotic minimaxity of wavelet estimators with sampled data.,” *Stat. Sin.*, 9(1), 1–32.
- GAIFFAS, S. (2005): “Sharp estimation in sup norm with random design,” Preprint 1010, LPMA Université Paris VI.
- GRAMA, I., AND M. NUSSBAUM (1998): “Asymptotic equivalence for nonparametric generalized linear models.,” *Probab. Theory Relat. Fields*, 111(2), 167–214.
- HOFFMANN, M., AND O. LEPSKI (2002): “Random rates in anisotropic regression. (With discussion).,” *Ann. Stat.*, 30(2), 325–396.
- JOHNSTONE, I. M., AND B. W. SILVERMAN (2004): “Boundary coiflets for wavelet shrinkage in function estimation.,” *J. Appl. Prob.*, Special Vol. 41A, 81–98.
- LE CAM, L., AND G. LO YANG (2000): *Asymptotics in statistics. Some basic concepts. 2nd ed.* Springer Series in Statistics. New York, Springer.
- MEYER, Y. (1995): *Wavelets and operators.* Cambridge University Press, Cambridge.
- NUSSBAUM, M. (1996): “Asymptotic equivalence of density estimation and Gaussian white noise.,” *Ann. Stat.*, 24(6), 2399–2430.
- ROHDE, A. (2004): “On the asymptotic equivalence and rate of convergence of nonparametric regression and Gaussian white noise.,” *Stat. Decis.*, 22(3), 235–243.
- SWELDENS, W., AND R. PIESSENS (1993): “Wavelet Sampling Techniques,” in *1993 Proceedings of the Statistical Computing Section*, pp. 20–29. American Statistical Association.
- WOJTASZCZYK, P. (1997): *A mathematical introduction to wavelets.* London Mathematical Society Student Texts. 37. Cambridge: Cambridge University Press.